


Weeds Watch Out!
Stop Invasive Aquatic Plants

# Aquatic Plants: The Good & The Bad

ur lakes and rivers would seem barren without lush plants along the shore. Plants do more than give us pleasing vistas—they are havens for wildlife and birds. Fish rely on aquatic plants for food and habitat. Plants improve water quality by absorbing nutrients and filtering pollutants. They protect the lake shoreline by holding soil on the lake bottom and water's edge—which in turn reduces erosion. Aquatic plants are an important part of the lakeshore environment.

As beneficial as aquatic plants are, some can create problems and become nuisance weeds. Some plants not native to an area are called "invasive" or "nuisance" plants. As they proliferate, invasive aquatic plants can impede boaters and swimmers, and generally lower the aesthetic and economic value of the waterbody. Infested waterways decrease property values, hurt tourism, impact fisheries, and cost communities money required to control and manage the invasive weeds.

Non-native, invasive species are introduced into and spread throughout the Finger Lakes region by various methods. One of the easiest, and most common ways for invasive weeds to be introduced is by "hitching a ride" on boats as they navigate various waterbodies. Some species can also be introduced unintentionally when they are used in gardens and



landscaping near a waterway. Additionally, when people dispose of aquatic plants by emptying their aquariums into a nearby waterway, non-native, nuisance plants are introduced into the region.

Preventing the introduction and spread of invasive weeds is essential to the health of the Finger Lakes region. You can help, starting with early detection. Learn to identify native and non-native plants. If invasive plants are spreading to a new area, report the sighting to the project leaders listed on the back cover of this guide.

# Glossary and Plant Index

### Glossary

Alternate: Leaves spaced singly along a stem

Emergent: Found in shallow water with a large portion of stems and leaves growing above (emerging from) the water surface

Floating: Have leaves that float on top of the water

Leaflets: Individual segments radiating from a common stem, that together make up a leaf

Lobe: Portion of a leaf that projects outward and divides the leaf into distinct parts, but not enough to make them separate leaflets; lobes may be rounded or pointed

Margin: Leaf edge

Mid-rib: Central vein of a leaf, running from base to tip

Opposite: Leaves spaced in pairs, one on each side of the stem

Rosette: Leaves arranged in a radiating pattern at the base of the plant

Submersed: Growing underwater

Toothed margin: Shallow bumps

Tuber: Thickened portion of stem, providing food storage for the plant

Whorl: Arrangement of 3 or more leaves or flowers radiating from a common point

This guide will help you to recognize the difference between aquatic invasive weeds and their native look-a-likes found in the Central New York region. In addition, there are suggestions as to what to do if you identify an invasive weed. For the purpose of this guide, plants are separated into two categories: Invasive Weed, on the left, and Native Plant, on the facing page. Note that several invasive weeds share one or two native look-a-likes.

Invasive Weed: Non-native nuisance plants that have been identified either in the Finger Lakes Region or in neighboring watersheds. It is important to prevent the spread of these plants.

Native Plant: Plants that look similar to invasive weeds but are beneficial to the ecosystem, and should be left alone and allowed to grow.



### Floating plants:

• F	ree floater—E	uropean frogbit	(p. 4)
-----	---------------	-----------------	--------

Rooted to sediment—Water chestnut (p. 6)




### Submersed plants:

<ul> <li>Whorl leaf pattern—Hydrilla</li> </ul>	(p. 8)
Whorl leaf pattern—Brazilian waterweed	(p. 10)

- Lasagna-like leaves—Curly-leaved pondweed (p. 12)
- Grass-like leaves—Starry stonewort (p. 14)
- Finely divided leaves—Eurasian watermilfoil (p. 16)
- Finely divided leaves—Fanwort (p. 18)
- Finely divided leaves—Parrotfeather (p. 20)

# Control & Eradication Techniques






Early detection and rapid response are important. To be sure of proper identification and early detection of invasive weeds, please use this guide to become familiar with both the native and non-native plants in the region. If you find an invasive weed in a new area within the Finger Lakes region, please inform one of the information contacts listed on the back of this guide at once so that new populations of invasive plants can be verified and promptly removed.

These plants both grow and spread by various methods, and therefore control and eradication planning need to be specific to each plant species. For instance, some plants spread by fragmentation. Plants that spread by fragmentation can grow from pieces of the original plant with no need for seeds. Other plants reproduce by seeds or buds. To maximize the benefit, control methods should be undertaken in the spring or very early summer before plants set seed.

Control techniques include physical, chemical and biological methods. Physical control includes raking, cutting, or harvesting vegetation. Since many invasive weeds can spread by fragmentation, care should be taken to prevent fragments from being carried away by water. In some cases, hand pulling can be an effective control technique, as long as the entire plant is pulled, not just the upper portion. Biological control uses specific insects or fish as natural predators of invasive weeds. Chemical methods require permits from the NYS Department of Environmental Conservation. Please contact one of the informational contacts on the back of this guide for more information.

Even with various control methods available, prevention is a vital first step, and you can help! Actions to take include cleaning boats and equipment on dry land, early detection, and quickly informing any leader listed on the back of the guide if you spot a new infestation of an invasive weed in the Finger Lakes region.







Opposite Leaf
Pattern

Plant posing a threat to the Finger Lakes Region

#### European frogbit (Hydrocharis morsus-ranae)



Beneficial plants that look like Frogbit and Water chestnut



### Assume Spatterdock if:


- Heart-shaped leaves up to 16 inches long.
- Parallel or overlapping rounded lobes.
- · Yellow flowers.
- Large mid-rib.
- · Found in ponds or slow moving waters.

#### Assume Water lily if:


- Round leaves with pointed lobes, 6–8 inches in length.
- Many-petalled white flowers float on water surface.
- Veins on leaf radiate out from where leaf and stem meet.
- Found in quiet waters.

Plant posing a threat to the Finger Lakes Region

#### Water chestnut (Trapa natans)



Beneficial plants that look like Water chestnut and Frogbit




### Assume Water lily if:

- Round leaves with pointed lobes, 6–8 inches in length.
- Many-petalled white flowers float on water surface.
- Veins on leaf radiate out from where leaf and stem meet.
- Found in quiet waters.

### Assume Spatterdock if:

- Heart-shaped leaves, up to 16 inches long.
- Parallel or overlapping rounded lobes.
- · Yellow flowers.
- Large mid-rib.
- Found in ponds or slow moving waters.


Plant posing a threat to the Finger Lakes Region



Beneficial plants that look like Hydrilla and Brazilian waterweed

Southern naiad (Najas guadalupensis) American elodea (Elodea, spp.)





#### Assume Southern naiad if:

- Leaves are narrow, ribbon-like with broad base where they attach to stem.
- Leaves are arranged in pairs on opposite sides of the stem.
- The leaf is tapered with small teeth.
- Leaves are deep-green to purple-green.


#### Assume American elodea if:

- Small, lance-shaped leaves in whorls of no more than three.
- Leaves appear and feel smooth.
- Lack of spines on the mid-rib.
- No tubers when pulled from the sediment.


Plant posing a threat to the Finger Lakes Region

### Brazilian waterweed (Egeria densa)

Plant is much "thicker" than Elodea.



Beneficial plants that look like Brazilian waterweed and Hydrilla



American elodea (Elodea spp.) Southern naiad (Najas guadalupensis)



#### Assume American elodea if:

- Small, lance-shaped leaves in whorls of no more than three.
- Leaves appear and feel smooth.
- Lack of spines on the mid-rib.
- No tubers when pulled from the sediment.

#### Assume Southern naiad if:

- Leaves are narrow, ribbon-like with broad base where they attach to stem.
- Leaves are arranged in pairs on opposite sides of the stem.
- The leaf is tapered with small teeth.
- Leaves are deep-green to purple-green.

Plant posing a threat to the Finger Lakes Region



Beneficial plant that looks like Curly-leaved pondweed


Clasping-leaved pondweed (Potamogeton perfoliatus)






- Assume Clasping-leaved if:
  - Wide, oval-shaped leaves have smooth edges.
- In shallow water, plants have darker foliage than plants growing in deeper water.
- Basal parts of leaves clasp straight, plant stems are slender.

Plant posing a threat to the Finger Lakes Region



Beneficial plants that look like Starry stonewort

Nitella (Nitella, spp.) Muskgrass (Chara, spp.)







- · Branches are smooth and flexible.
- Branches/leaves are translucent green.
- Branches arranged in whorls around stem.
- Lacks roots. Attaches to the sediment by root-like structures called "holdfasts."
- May be free-floating above sediment.

### Assume Muskgrass if:

- Plant is covered in a brittle, scaly coating.
- · Often smells "skunky" when pressed.
- Branches are hard and ridged.
- · Feels gritty when crushed.
- Inhabits fresh to brackish (salty) waters; both shallow and deep.


Plant posing a threat to the Finger Lakes Region

### Eurasian watermilfoil (Myriophyllum spicatum)



Beneficial plants that look like Eurasian watermilfoil

Northern watermilfoil (Myriophyllum sibiricum)
Coontail (Ceratophyllum demersum)



#### Assume N. watermilfoil if:


- Whorls of 4~5 of feather-like leaves.
- Leaves made up of 5-12 thread-like leaflets.
- Leaves tend to be stiff when removed from water.
- Leaf tip is tapered.
- Tips and shoots lack reddish-brown color.
- · Inhabits lakes, ponds, and rivers.

#### Assume Coontail if:


- Leaves are not feather-like.
- Bristle-like leaves are toothed and have forked division.
- Whorls of leaves are tight at tips, resembling a raccoon tail.
- Leaves keep their shape out of water.
- Inhabits slow moving waters of streams and rivers, as well as lakes and ponds.

Plant posing a threat to the Finger Lakes Region

#### Fanwort (Cabomba caroliniana)



Beneficial plant that looks like Fanwort.




stem.

Produces yellow flowers July to September.Can be found on lakeshores and in ponds

and slow moving streams.

Plant posing a threat to the Finger Lakes Region

### Parrotfeather (Myriophyllum aquaticum)



- Leaves are rigid, deeply serrated, and profuse.
- Woody stems can grow over 5 feet long.
- Prefers slow-moving waters of streams, rivers and ditches; also found in shallow lakes and ponds.

Weeds Watch Out! is an education and outreach program that will help to control the spread of invasive aquatic plant species into, within, and from the Finger Lakes Region. W2O! has established strong inter-watershed relationships in the Finger Lakes Region to effectively address current and future invasive aquatic plant issues in a coordinated manner. For more information, visit the web at www.co.cayuga.ny.us/wqma/weedswatchout http://watershed.extendonondaga.org www.nyis.info

W2O! was funded by the U.S. Environmental Protection Agency (EPA) Great Lakes Grants Program, E.P.A. Great Lakes National Program Office, the National Fish and Wildlife Foundation and the Finger Lakes Institute. The views and conclusions contained in this document are those of the authors and should not be interpreted as representing the opinions or policies of the U.S. Government or the National Fish and Wildlife Foundation. Mention of trade names or commercial products does not constitute their endorsement by the U.S. Government or the National Fish and Wildlife Foundation.

Reprinted with modifications in 2014 with funding from the Finger Lakes Institute and City of Syracuse Water Department.

Line drawings: University of Florida, Center for Aquatic and Invasive Plants Line Drawing DVD Master; USDA, NRCS. 2004. The PLANTS Database, Version 3.5 (http://plants.usda.gov). Used with permission.

Photographs: Robert Johnson, Cornell University. Ruthanna Hawkins, Cayuga Lake Watershed Network. Used with permission. ©CLWN 2005 10M alr.indd

#### **Information Contacts:**



# Cornell University Cooperative Extension Cayuga and Onondaga Counties

Cayuga County cayuga@cornell.edu 315.255.1183

Onondaga County onondaga@cornell.edu 315.424.9485



City of Syracuse Water Department 315.685.6486





Cayuga Lake Watershed Network steward@cayugalake.org 607.532.4104



Finger Lakes Institute fli@hws.edu 315.781.4390



Oswego County Soil & Water information@oswegosoilandwater.com 315.592.9663



Cayuga County Planning & Development wqma@co.cayuga.ny.us 315.253.1276